

The Ergogenics of Hypoxia Training in Athletes

Brett M. Loffredo, MD, and James L. Glazer, MD

Corresponding author

James L. Glazer, MD
Division of Sports Medicine, Maine Medical Center,
Department of Family Medicine, 272 Congress Street,
Portland, ME 04101, USA.
E-mail: glazej@mmc.org

Current Sports Medicine Reports 2006, 5:203–209
Current Science Inc. ISSN 1537-890x
Copyright © 2006 by Current Science Inc.

Hypoxia elicits hematopoiesis, which ultimately improves oxygen transport to peripheral tissues. In part because of this, altitude training has been used in the conditioning of elite endurance athletes for decades, despite equivocal evidence that such training benefits subsequent sea level performance. Recently, traditional live high-train high athletic conditioning has been implicated in a number of deleterious effects on training intensity, cardiac output, muscle composition, and fluid and metabolite balance—effects that largely offset hematopoietic benefits during sea level performance. Modified live high-train low conditioning regimens appear to capture the beneficial hematopoietic effects of hypoxic training while avoiding many of the deleterious effects of training at altitude. Because of the logistical and financial barriers to living high and training low, various methods to simulate hypoxia have been developed and studied. The data from these studies suggest a threshold requirement for hypoxic exposure to meaningfully augment hematopoiesis, and presumably improve athletic performance.

Introduction

There is little debate that training at altitude improves athletic performance at altitude [1]. For decades, elite athletes have invested considerable resources in training at altitude, with the intention of improving performance at sea level. Studies indicate that there are numerous physiologic changes associated with hypoxic and hypobaric training—some advantageous and some detrimental to athletic performance at or near sea level. Recent advances have allowed athletes to optimize the beneficial effects of hypoxic/hypobaric training while minimizing deleterious effects. Training modalities such as live high-train

low (LHTL), intermittent hypoxic exposure (IHE), hypoxic/hypobaric chambers, and exercise hypoxia have changed the face of state of the art training.

This review, although by no means comprehensive, looks at the effects of hypoxic/hypobaric training from a performance enhancement and systems-based perspective. Training recommendations are made, and future research is suggested.

Performance-based Responses to Altitude

Altitude training has been incorporated into endurance athletes' preparation for decades. At first, athletes employed a traditional live high-train high (LHTH) exercise program. Research failed to show that this improves aerobic performance at sea level, although anaerobic performance may be enhanced [2–4]. More recently, modified altitude training regimens (ie, LHTL) have been shown in some well-designed studies to improve subsequent endurance performance at sea level [5,6,7,8,9]. Because altitude training, especially LHTL, is logistically difficult as well as financially restrictive for many athletes, newer hypoxic training modalities have been developed, including intermittent hypoxic exposure, hypoxic tents, and hypoxic exercise. Studies of such simulated altitude exposure have produced mixed results with regard to performance [10].

Altitude training appears to benefit elite athletes principally through an improvement in oxygen transportation. This is achieved largely by an increase in hematocrit and blood volume [11,12], although a concomitant volume contraction and decrements in cardiac output in short-term altitude exposure may mitigate these benefits [13,14]. Other deleterious effects of exercise at altitude include a relative detraining effect, which can offset any improvements in oxygen-carrying capacity [15,16].

Optimal endurance performance relies upon frequency, duration, and intensity of training [17,18]. Especially with endurance performance [19], maintenance of training intensity appears to be the principle variable in optimizing subsequent endurance performance [20,21]. Failure to maintain training intensity results in lower VO_2max [20–24], and a subsequent detraining effect [15,16]. The reasons for the detraining effect appear to be multifactorial. Oxygen uptake at a main-

tained power output at sea level and altitude is similar [6,25–27]. Because VO_2max declines with increasing altitude (a function of decreased atmospheric oxygen content), a given power output at altitude will correspond with a greater relative exercise intensity (higher % VO_2max). At the same absolute intensity, exercising at altitude results in higher blood lactate, heart rate, ventilation, and perceived effort compared with similar training at sea level [13,28]. Athletes training at altitude therefore self-select reduced work rates [5,6]. Thus, training power output is typically reduced at altitude compared with sea level for any given amount of work done [5,29], with subsequent relative deconditioning in altitude training versus sea-level training [15,16].

LHTL programs and simulated intermittent hypoxia/hypobaric training regimens have been designed to capture the beneficial hematopoietic effects of altitude exposure, while avoiding the detraining effects of altitude. Results of several well-designed studies involving LHTL have proven benefits to subsequent sea-level performance in endurance athletes [6,30••]. The performance enhancement during these LHTL studies can be impressive. For example, over a 27-day training period, elite runners showed a 1.1% improvement in 3000 meter time trial time in spite of their recent participation in a national championship competition [30••]. One third of the athletes achieved a personal best time subsequent to the LHTL training protocol. Results were similar among both male and female participants.

IHE can avoid the intrinsic logistical and financial constraints of LHTL. Some studies of IHE have also investigated the threshold hypoxic stimulus required to effect performance enhancement. Initial uncontrolled studies of IHE reported significant improvements in anaerobic threshold, hematologic measures, and ventilatory measures after 17 days of 3 to 5 hours per day of altitude exposure at 4000 to 5500 meters [31]. The same research group had previously demonstrated no difference in performance benefit whether the hypoxia was administered at rest or during exercise [32].

Evidence for performance enhancement in simulated hypoxic environments can be extrapolated from data regarding hematopoiesis. Some studies have found no erythropoietic effect with normobaric hypoxia during sleep [33–35], although hypoxic exposures were for less than 10 hours per day for less than 3 weeks. Other researchers have found an increase in erythrocyte mass with exposures of 16 hours per night for 4 weeks [36,37]. Studies of 12 hours per day of hypoxia during rest at a simulated altitude of 2000 to 2700 meters failed to show an improvement in VO_2 , although erythropoietin and reticulocyte counts were improved [38]. Significant improvement in 3000-meter timed trial after 3 hours per day of hypoxia for 14 days was reported [39]. A more recent study of 70 minutes a day of alternate 5-minute cycles of hypoxia and normoxia 5 days a week for

4 weeks did not show improvement in hematologic or performance measures [40]. These studies together suggest a threshold dose of hypoxia required for augmented hematopoiesis [10•]. The details regarding the degree and duration of hypoxia required for athletic enhancement remain to be elucidated.

Putative Mechanisms of Physiologic Adaptation During Altitude Training Hematologic

Altitude exposure augments the hematopoietic response. This has been postulated as one of the key benefits of altitude training; transfusion studies and recombinant erythropoietin studies suggest that increased erythrocyte mass alone increases VO_2max for endurance athletes regardless of performance level [41–45]. There are also deleterious hematologic effects associated with brief altitude exposure, including volume contraction and subsequent perfusion decrements. Training modalities such as LHTL may maximize the erythropoietic effects of altitude training while concomitantly reducing deleterious hemodynamic effects.

Natives living above 2500 meters demonstrate a significantly higher hemoglobin, hematocrit, and erythrocyte volume compared with sea level residents [12,46]. The polycythemia of altitude residence seems to depend on the duration of habitation at altitude [47,48]. Studies suggest a threshold altitude of 2200 to 2500 meters, corresponding to a threshold PO_2 of 70 mm, is needed for altitude-dependent hematopoiesis [12]. The hematopoietic response is greater at higher altitudes, likely secondary to increasing hypoxia and subsequent release of higher levels of erythropoietic factors at higher altitude [12].

Hematopoiesis increases continually during altitude training. Berglund [48] found that hemoglobin concentration during acute altitude training increased at approximately 1% per week. The apparent benefit to athletes of increased hemoglobin concentration may be offset by concomitant sympathetically mediated arterioconstriction and decreased cardiac output, which combine to prevent an increase in O_2 transport overall for those undergoing traditional altitude training [14]. In addition to those cardiovascular effects, athletes during an altitude sojourn may lose up to 25% of their plasma volume [13], although athletes who reside at altitude appear to maintain plasma volumes comparable with athletes residing at sea level [49]. Such hemodynamic changes may contribute to a decrement in muscle perfusion at altitude which persists upon return to sea level [14].

LHTL has been shown to increase markers of hematopoiesis, erythrocyte volume, and hemoglobin concentration in competitive and elite athletes [6,30••]. These findings have been correlated with increased VO_2max and performance measures [6,30••].

The results of studies of simulated altitude exposure have been mixed with regard to hematopoietic effects.

Twenty-three nights of 8 to 10 hours at a simulated altitude of 3000 meters in a nitrogen house did not elicit a change in total hemoglobin mass or hematopoietic markers compared with control subjects [33]. A 4-week regimen of alternating 5-minute cycles of hypoxia and normoxia for 70 minutes five times a week did not alter erythropoietic markers in highly trained runners versus control subjects [40]. Among well-trained runners, an 8- to 11-hour per night exposure to a simulated altitude of 2650 meters for 5 nights, followed by a 3-night sojourn, and then repeated for three cycles, resulted in an increase in serum erythropoietin but did not increase reticulocyte production versus control [35]. These findings are consistent with the suggestion that there are minimum requirements for the degree and duration of hypoxia required for effective erythropoiesis.

Respiratory

Altitude training has both positive and negative effects on the respiratory conditioning of athletes. Training at altitude may result in improved respiratory muscle training because of an increase in resting and exercise ventilation [50]. This increase in ventilatory exchange (VE) at altitude is well documented [51–53], and has been associated with increased respiratory muscle work at altitude, which persists upon return to sea level [52]. The primary mechanism of increased VE at altitude is augmentation of the hypoxic ventilatory response (HVR), which has been found during continuous altitude residence [54], intermittent hypoxic exposure [55], and LHTL [56,57]. The increased HVR facilitates a progressive rise in VE despite unchanged hypoxic stimulus and hypercapnic inhibition [51]. On return to normoxia, this hyperventilation continues and then diminishes in a time-dependent manner [51].

The physiologic effects of this increased ventilatory exchange during exercise are unclear. Hyperpnea during exercise may be detrimental because of reduced peripheral blood flow to compensate for the greater blood flow demands of breathing [58]. Conversely, insufficient exercise hyperpnea may lead to exercise-induced hypoxemia at submaximal workloads [59,60], although it is unclear whether this affects performance.

Traditional altitude training has been found to augment HVR and VE after 2 to 4 weeks [61–64]. The duration and degree of hypoxia required to augment HVR and VE in subsequent sea level performance using an LHTL protocol or simulated altitude exposure is unclear. As little as 1 hour a day of hypoxia for 7 days was enough to increase HVR, although submaximal VE remained unchanged [65]. Another study found that 45 minutes a day of 2500-meter hypoxia, 5 days a week for 5 weeks during exercise increased HVR but not VE [5]. Subjects exposed to hypoxia for 8 to 10 hours a night at 3000 meters showed increased submaximal exercise VE in normoxia after 11 days, although HVR was not measured [57]. Other research has found a

37% increase in ventilation during cycling at fixed velocity after 8 days of LHTL [66].

A more recent controlled study found that exposure to 4 nights of LHTL increased submaximal exercise VE measured in normoxia by 9%, and remained elevated throughout a 3-week intervention [67]. This increased VE was strongly correlated with an increased HVR by 15 to 19 days of LHTL, with a maximal HVR at 19 days of training—effects that were not noted in sea level control subjects [67]. This is the first study to report increased submaximal VE in normoxia after 4 days of LHTL in well-trained athletes [67].

No increased $\text{VO}_{2\text{max}}$ was noted after 19 days of LHTL [67]. This finding is in accord with most previous studies of traditional altitude training and LHTL alike, which raises the question whether increased submaximal but not maximal VE is a reflection that trained athletes may be limited by mechanical constraints of the pulmonary system at high workloads [68]. It has been suggested that expiratory flow limitation during maximal exercise limits $\text{VO}_{2\text{max}}$ [69,70].

$\text{VO}_{2\text{max}}$ during acute altitude exposure in athletes is diminished, even at altitudes as low as 580 meters [71] and 900 meters [72]. The decrement in $\text{VO}_{2\text{max}}$ in hypoxia is directly proportional to $\text{VO}_{2\text{max}}$ in normoxia, suggesting that athletes are affected more negatively by altitude exposure than untrained subjects. It is unclear whether observed decrements in $\text{VO}_{2\text{max}}$ affect performance, as significant improvement in exercise performance without improved $\text{VO}_{2\text{max}}$ has been demonstrated [1,18,63,73–75]. Furthermore, improvements in $\text{VO}_{2\text{max}}$ may not necessarily evoke improvement in exercise performance [6].

Cardiovascular

The effects of traditional altitude training on the cardiovascular system appear to be largely deleterious to sea level performance. Although chronic exposure to hypobaric hypoxia increases arterial oxygen concentration as a result of an increased hemoglobin concentration, sympathetically mediated arterioconstriction and decreased cardiac output prevent an increase in oxygen transport [14]. The proposed mechanism for the reduction in cardiac output observed at altitude is decreased vegetative stimulation [13], possibly by downregulation of β -adrenoceptors [76]. This may lead to a negative altitude training effect on the heart. It is unclear whether modified altitude training regimens, such as LHTL and IHE, avoid the cardiac detraining associated with chronic altitude exposure.

Musculoskeletal

The effects of prolonged altitude exposure on muscle structure and function appear to be largely deleterious. Prolonged exposure to altitudes above 4500 meters has been shown to result in reduced muscle mass [77]. Muscle perfusion is decreased at altitude, and that perfusion deficit persists after return to sea level [14,78]. Above 4300 meters, altitude exposure is associated with loss of mitochondria and oxidative enzyme capacity [79]. Simi-

Table 1. Quality of evidence: ergogenics of hypoxia training

Study	Study findings	Strength of evidence
Martino et al. [3], Nummela et al. [4], Levine and Stray-Gundersen [6], Levine et al. [7•], Levine et al. [8], Levine et al. [9], Stray-Gundersen et al. [30••]	Live high-train low improves athletic performance	Level 1
Katayama et al. [39]	Simulated hypoxic exposure improves athletic performance	Level 2
None listed	Live high-train high improves sea level athletic performance	Level 3
Boutellier et al. [14], Gore et al. [71]	Live high-train low harms sea level athletic performance	Level 2

Table 2. Assessing quality of evidence

Study quality	Diagnosis	Treatment, prevention, screening, outcomes	Prognosis
Level 1: good quality patient oriented evidence	Validated clinical decision rule	SR/meta-analyses or randomized controlled trials with consistent findings	SR/meta-analysis of good-quality cohort study
	SR/meta-analyses of high-quality studies	High-quality individual randomized controlled trials	Prospective cohort study with good follow-up
	High-quality diagnostic cohort study	All or none study	
Level 2: limited quality patient-oriented evidence	Unvalidated clinical decision rule	SR/Meta-analysis of lower-quality clinical trials or of studies with inconsistent findings	SR/Meta-analysis of lower quality cohort studies or with inconsistent results
	SR/meta-analysis of lower-quality studies or studies with inconsistent findings	Lower quality clinical trial	Retrospective cohort study or prospective cohort study with poor follow-up
	Lower quality diagnostic cohort study or diagnostic case-control study	Cohort study	Case-control study
		Case-control study	Case series
Level 3: other evidence	Consensus guidelines, extrapolations from bench research, usual practice, opinion, disease-oriented evidence (intermediate or physiologic outcomes only), or case series for studies of diagnosis, treatment, prevention, or screening.		

SR—systematic review

(Adapted from Ebell et al. *Am Fam Physician* 2004, 69:549–557.)

lar results have been documented after altitude training at 2700 meters, and low-pressure chamber training with equal relative intensity [2,80,81]. Athletes exposed to an equal absolute workload showed increased oxidative capacity and myoglobin concentration on return to normoxia [79,81–83]. Other studies indicate that anaerobic metabolic enzymes are not enduringly affected by altitude exposure [84]. The effects of hypoxia on muscle capillary

networks is unclear [2,80,81,85]. Only Mizuno et al. [2] and Hoppeler and Vogt [86] documented increased muscle capillarity after altitude exposure, possibly due to a confounding effect of improved conditioning in muscles not fully trained previously.

Chronic altitude residence has previously been found to augment muscle oxidative capacity [87]. Muscle biopsies have shown higher myoglobin and oxidative enzyme

concentrations in high altitude natives compared with sea level counterparts [87]. These initial findings may simply be due to differences in training level between altitude and sea level subjects [88,89].

Conclusions

Training at altitude improves performance at altitude. Increasingly, data suggest that traditional LHTH conditioning is deleterious to subsequent sea level performance. Intermittent hypoxic exposure, as best exemplified by several studies of LHTL training regimens, has been shown to improve sea level performance. Simulated hypoxic exposure, although promising, has shown mixed results. The lack of a consistent improvement in performance or markers of hematopoiesis during studies of simulated hypoxic exposure is likely secondary to the great variability in duration of hypoxic exposure among such studies. These studies suggest a threshold degree/duration of hypoxia required for significant hematopoiesis and subsequent performance enhancement. As little as 3 hours of hypoxia a day for 2 weeks has been linked to improvement in subsequent timed trial performance [39]. Other studies of IHE with various training parameters have shown improved hematologic measures [36–38], although others have shown no such improvement [33–35].

Based on a review of current literature, only LHTL conditioning has been shown to improve performance in well-controlled prospective studies. There is no evidence that hypoxia during training benefits performance. In fact, studies of traditional altitude training suggest that training in hypoxia has the negative effect of reduced training intensity. Simulated hypoxic exposure studies are promising, but have yet to be proven to augment athletic performance definitively (Tables 1 and 2).

References and Recommended Reading

Papers of particular interest, published recently, have been highlighted as:

- Of importance
- Of major importance

1. Maher JT, Jones LG, Hartley LH: **Effects of high-altitude exposure on submaximal endurance capacity of men.** *J Appl Physiol* 1974, 37:895–898.
 2. Mizuno M, Juel C, Bro-Rasmussen T, et al.: **Limb skeletal muscle adaptation in athletes after training at altitude.** *J Appl Physiol* 1990, 68:469–502.
 3. Martino M, Myers K, Bishop P: **Effects of 21 days training at altitude on sea-level anaerobic performance in competitive swimmers.** *Med Sci Sports Exerc* 1996, 27(Suppl 5):abstract 37.
 4. Nummela A, Joutte P, Rusko H: **Effect of living high and training low on sea level anaerobic performance in runners.** *Med Sci Sports Exerc* 1996, 28(Suppl 5):abstract 740.
 5. Levine BD, Stray-Gundersen J: **A practical approach to altitude training: where to live and train for optimal performance enhancement.** *Int J Sports Med* 1992, 13(Suppl 1):S209–S212.
 6. Levine BD, Stray-Gundersen J: **Living high-training low: effect of moderate-altitude acclimatization with low-altitude training on performance.** *J Appl Physiol* 1997, 83:102–112.
 7. Levine BD, Friedmann B, Stray-Gundersen J: **Confirmation of the “high-low” hypothesis: living at altitude-training near sea level improves sea level performance.** *Med Sci Sports Exerc* 1996, 28:abstract 742.
- The first well-controlled study documenting improved sea level athletic performance after a LHTL training protocol.
8. Levine BD, Stray-Gundersen J, Duhaime G, et al.: **Living high-training low: the effect of altitude acclimatization/normoxic training in trained runners.** *Med Sci Sports Exerc* 1991, 23:abstract 145.
 9. Levine BD, Engfred K, Friedman DB, et al.: **High altitude endurance training: effect on aerobic capacity and work performance.** *Med Sci Sports Exerc* 1990, 22:abstract 209.
 10. Levine BD: **Intermittent hypoxic training: fact and fancy.** *High Alt Med and Biol* 2002, 3:177–193.
- A contemporary review of recent developments in the study of intermittent hypoxic exposure.
11. Hurtado A, Merino C, Delgado E: **Influence of anoxemia on the hematopoietic activity.** *Arch Intern Med* 1945, 75:284–323.
 12. Weil JV, Jamieson G, Brown DW, Grover RF: **The red cell mass – arterial oxygen relationship in normal man.** *J Clinical Invest* 1968, 47:1627–1639.
 13. Young AJ, Young PM: **Human acclimatization to high terrestrial altitude.** *Human Performance Physiology and Environmental Medicine at Terrestrial Extremes.* Indianapolis: Benchmark Press; 1988:497–543.
 14. Boutellier U, Marconi C, DiPrampero PE, Cerretelli P: **Effects of chronic hypoxia on maximal performance.** *Clin Respir Physiol* 1982, 18(Suppl 4):39–44.
 15. Balke B: **Work capacity and its limiting factors at high altitude.** Weihe WH, ed. *The physiological effects of high altitude.* New York: Pergamon Press; 1964:233–247.
 16. Rusco HK, Kirvesniemi H, Paavola L: **Effect of altitude training on sea level aerobic and anaerobic power of elite athletes [abstract].** *Med Sci Sports Exerc* 1996, 28:S124.
 17. Hickson RC, Bomze HA, Holloszy JO: **Linear increase in aerobic power induced by a strenuous program of endurance exercise.** *J Appl Physiol* 1977, 42:372–376.
 18. Pollock ML: **The quantification of endurance training programs.** Wilmore JH, ed. *Exercise and sport science reviews.* New York: Academic Press; 1973:155–188.
 19. McConnell GK, Costill DL, Widrick JJ: **Reduced training volume and intensity maintain aerobic capacity but not performance in distance runners.** *Int J Sports Med* 1993, 14:33–37.
 20. Hickson RC, Foster C, Pollock ML: **Reduced training intensities and loss of aerobic power, endurance, and cardiac growth.** *J Appl Physiol* 1985, 58:492–499.
 21. Hickson RC, Kanakis C, Davis JR: **Reduced training duration and effects on aerobic power, endurance, and cardiac growth.** *J Appl Physiol* 1982, 53:225–229.
 22. Hickson RC, Rosenkoetter MA: **Reduced training frequencies and maintenance of increased aerobic power.** *Med Sci Sports Exerc* 1981, 13:13–16.
 23. McConnell GK, Costill DL, Widrick JJ: **Reduced training volume and intensity maintain aerobic capacity but not performance in distance runners.** *Int J Sports Med* 1993, 14:33–37.
 24. Pollock ML: **The quantification of endurance training programs.** In *Exercise and Sport Science Reviews.* Edited by Wilmore JH. New York: Academic Press; 1973:155–188.
 25. Fulco CS, Lewis SF, Frykman P: **Muscle fatigue and exhaustion during dynamic leg exercise in normoxia and hypobaric hypoxia.** *J Appl Physiol* 1996, 81:1891–1900.
 26. Hansen JE, Vogel JA, Stelter GP, Consolazio CF: **Oxygen uptake in man during exhaustive work at sea level and high altitude.** *J Appl Physiol* 1967, 23:511–522.
 27. Knuttgen HG, Saltin B: **Oxygen uptake, muscle high-energy phosphates, and lactate in exercise under acute hypoxic conditions in man.** *Acta Physiol Scand* 1973, 87:368–376.

28. Sutton JR: **Effect of acute hypoxia on the hormonal response to exercise.** *J Appl Physiol* 1977, 42:587–592.
29. Fulco CS, Rock PB, Cymerman A: **Maximal and submaximal exercise performance at altitude.** *Aviat Space Environ Med* 1998, 69:793–801.
- 30.●● Stray-Gundersen J, Chapman RF, Levine BD: **“Living high-training low” altitude training improves sea level performance in male and female elite runners.** *J Appl Physiol* 2001, 91:1113–1120.
- Definitive prospective controlled study of LHTL and athletic performance in elite long-distance runners
31. Casas M, Casas H, Pages T, et al.: **Intermittent hypobaric hypoxia induces altitude acclimation and improves the lactate threshold.** *Aviat Space Environ Med* 2000, 71:125–130.
32. Rodriguez FA, Casas H, Casas M, et al.: **Intermittent hypobaric hypoxia stimulates erythropoiesis and improves aerobic capacity.** *Med Sci Sports Exerc* 1999, 31:264–268.
33. Ashenden MJ, Gore CJ, Dobson GP, Hahn AG: **“Live high, train low” does not change the total haemoglobin mass of male endurance athletes sleeping at a simulated altitude of 3000m for 23 nights.** *Eur J Appl Physiol* 1999, 80:479–484.
34. Ashenden MJ, Gore CJ, Martin DT, et al.: **Effects of a 12 day “live high, train low” camp on reticulocyte production and haemoglobin mass in elite female road cyclists.** *Eur J Appl Physiol* 1999, 80:472–478.
35. Ashenden MJ, Gore CJ, Dobson GP, et al.: **Simulated moderate altitude elevates serum erythropoietin but does not increase reticulocyte production in well-trained runners.** *Eur J P Appl Physiol* 2000, 81:428–435.
36. Laitinen H, Alopaeus K, Heikkinen R, et al.: **Acclimatization to living in normobaric hypoxia and training in normoxia at sea level in runners [abstract].** *Med Sci Sports Exerc* 1995, 27:S617.
37. Rusko HK, Tikkanen H, Paavolainen L, et al.: **Effect of living in hypoxia and training in normoxia on sea level VO₂Max and red cell mass.** *Med Sci Sports Exerc* 1999, 31:S86.
38. Piehl-Aulin K, Svedenhag J, Wide L, et al.: **Short-term intermittent normobaric hypoxia – haematological, physiological, and mental effects.** *Scand J Med Sci Sports* 1998, 8:132–137.
39. Katayama K, Matsuo H, Ishida K, et al.: **Intermittent hypoxia improves endurance performance and exercise efficiency.** *High Alt Med Biol* 2003, 4:291–304.
40. Julian JG, Gore CJ, Wilber RL, et al.: **Intermittent normobaric hypoxia does not alter performance or erythropoietic markers in highly trained distance runners.** *J Appl Physiol* 2004, 96:1800–1807.
41. Buick FJ, Gledhill N, Froese AB, et al.: **Effect of induced erythrocythemia on aerobic work capacity.** *J Appl Physiol* 1998, 85:636–642.
42. Ekblom B, Goldberg AN, Gullbring B: **Response to exercise after blood loss and reinfusion.** *J Appl Physiol* 1972, 33:175–180.
43. Berglund B, Ekblom B: **Effect of recombinant human erythropoietin treatment on blood pressure and some haematological parameters in healthy men.** *J Intern Med* 1991, 229:125–130.
44. Birkeland, KI, Stray-Gundersen J, Hemmersbach P, et al.: **Effect of rhEPO administration on serum levels of sTfR and cycling performance.** *Med Sci Sports Exerc* 2000, 32:1238–1243.
45. Williams MH, Wesseldine S, Somma T, Schuster R: **The effect of induced erythrocythemia on 5-mile treadmill run time.** *Med Sci Sports Exerc* 1981, 13:169–175.
46. Sanchez C, Merino C, Figallo M: **Simultaneous measurement of plasma volume and cell mass in polycythemia of high altitude.** *J Appl Physiol* 1970, 28:775–778.
47. Schmidt W, Spielvogel H, Eckhardt KU, et al.: **Effects of chronic hypoxia and exercise on plasma erythropoietin level in natives living permanently at high altitude.** *J Appl Physiol* 1993, 74:1874–1878.
48. Berglund B: **High-altitude training. Aspects of haematological adaptation.** *Sports Med* 1992, 14:289–303.
49. Schmidt W, Heinicke K, Rojas J, et al.: **Blood volume and hemoglobin mass in endurance athletes from moderate altitude.** *Med Sci Sports Exerc* 2002, 34:1934–1940.
50. McMahon ME, Boutellier U, Smith RM, Spengler CM: **Hyperpnea training attenuates peripheral chemosensitivity and improves cycling endurance.** *J Exp Biol* 2002, 205:3937–3943.
51. Bisgard GE, Forster HV: **Ventilatory responses to acute and chronic hypoxia.** In *Handbook of Physiology: Environmental Physiology*. Edited by Fregly MJ, Blatteis CM. New York: Oxford University Press; 1996:1207–1239.
52. Levine BD, Friedman DB, Engfred K, et al.: **The effect of normoxic or hypobaric hypoxic endurance training on the hypoxic ventilatory response.** *Med Sci Sports Exerc* 1992, 24:769–775.
53. Steinacker JM, Halder A, Liu Y, et al.: **Hypoxic ventilatory response during rest and exercise after a Himalaya Expedition.** *Eur J Appl Physiol* 1996, 73:202–209.
54. Sato M, Severinghaus JW, Powell FL, et al.: **Augmented hypoxic ventilatory response in men at altitude.** *J Appl Physiol* 1992, 73:101–107.
55. Garcia N, Hopkins SR, Powell FL: **Effects of intermittent hypoxia on the isocapnic hypoxic ventilatory response and erythropoiesis in humans.** *Respir Physiol* 2000, 123:39–49.
- 56.● Townsend NE, Gore CJ, Hahn AG, et al.: **Living high-training low increases hypoxic ventilatory response of well-trained endurance athletes.** *J Appl Physiol* 2002, 93:1498–1505.
- A well-designed study with a good synopsis of current literature.
57. Gore CJ, Hahn AG, Aughey RJ, et al.: **Live high:train low increases muscle buffer capacity and submaximal cycling efficiency.** *Acta Physiol Scand* 2001, 173:275–286.
58. Wetter TJ, Harms CA, Nelson WB, et al.: **Influence of respiratory muscle work on VO₂ and leg blood flow during submaximal exercise.** *J Appl Physiol* 1999, 87:643–651.
59. Rice AJ, Scroop GC, Gore CJ, et al.: **Exercise-induced hypoxemia in highly trained cyclists at 40% peak oxygen uptake.** *Eur J Appl Physiol* 1999, 79:353–359.
60. Durand F, Mucci P, Prefaut C: **Evidence for an inadequate hyperventilation inducing arterial hypoxemia at submaximal exercise in all highly trained endurance athletes.** *Med Sci Sports Exerc* 2000, 32:926–932.
61. Buskirk ER, Kollias J, Akers RF, et al.: **Maximal performance at altitude and on return from altitude in conditioned runners.** *J Appl Physiol* 1967, 23:259–266.
62. Faulkner JA, Daniels JT, Balke B: **Effects of training at moderate altitude on physical performance capacity.** *J Appl Physiol* 1967, 23:85–89.
63. Daniels J, Oleridge N: **The effects of alternate exposure to altitude and sea level on world-class middle-distance runners.** *Med Sci Sports* 1970, 2:107–112.
64. Dill DB, Adams WC: **Maximal oxygen uptake at sea level and at 3090m altitude in high school champion runners.** *J Appl Physiol* 1971, 30:854–859.
65. Katayama K, Sato Y, Shima N, et al.: **Enhanced chemosensitivity after intermittent hypoxic exposure does not affect exercise ventilation at sea level.** *Eur J Appl Physiol* 2002, 87:187–191.
66. Mattila V, Rusko H: **Effect of living high and training low on the sea level performance in cyclists.** *Med Sci Sports Exerc* 1996, 28:S156.
67. Townsend NE, Gore CJ, Hahn AG, et al.: **Hypoxic ventilatory response is correlated with increased submaximal exercise ventilation after live high, train low.** *Eur J Appl Physiol* 2005, 94:207–215.
68. Dempsey JB: **JB Wolfe Memorial Lecture: is the lung built for exercise?** *Med Sci Sports Exerc* 1986, 18:143–155.
69. Dempsey JB, Hanson PG, Henderson KS: **Exercise-induced arterial hypoxaemia in healthy human subjects at sea level.** *J Physiol (Lond)* 1984, 355:161–175.

70. Johnson BD, Saupe KW, Dempsey JA: **Mechanical constraints on exercise hyperpnea in endurance athletes.** *J Appl Physiol* 1992, 73:874–886.
71. Gore CJ, Little SC, Hahn AG: **Reduced performance of male and female athletes at 580m altitude.** *Eur J Appl Physiol* 1997, 75:136–143.
72. Terrados N, Mizono M, Andersen H: **Reduction in maximal oxygen uptake at low altitudes: role of training status and lung function.** *Clin Physiol* 1985, 5(Suppl 3):75–79.
73. Astrand PO, Rodahl K: **Textbook of work physiology.** In *Physiological Bases of Exercise*, edn 3. New York: McGraw-Hill; 1986.
74. Gleser MA, Vogel JA: **Endurance capacity for prolonged exercise on the bicycle ergometer.** *J Appl Physiol* 1973, 34:438–442.
75. Horstman D, Weiskopf R, Jackson RE: **Work capacity during 3-wk sojourn at 4,300m: effects of relative polycythemia.** *J Appl Physiol* 1980, 49:311–18.
76. Antezana AM, Richalet J, Antezana G, et al.: **Adrenergic system in high altitude residents.** *Int J Sports Med* 1992, 13:S96–100.
77. Saltin B: **Aerobic and anaerobic work capacity at 2,300 metres.** *Med Thorac* 1967, 24:205–210.
78. Wolfel EE, Groves BM, Brooks GA, et al.: **Oxygen transport during steady-state submaximal exercise in chronic hypoxia.** *J Appl Physiol* 1996, 70:1129–1136.
79. Hoppeler H, Desplanches D: **Muscle structural modifications in hypoxia.** *Int J Sports Med* 1992, 13:S166–S168.
80. Saltin B, Kim CK, Terrados N, et al.: **Morphology, enzyme activities and buffer capacity in leg muscles of Kenyan and Scandanavian runners.** *Scan J Med Sci Sports* 1995, 5:222–230.
81. Terrados N, Melichna J, Sylven C, et al.: **Effects of training at simulated altitude on performance and muscle metabolic activity in competitive road cyclists.** *Eur J Appl Physiol* 1988, 57:203–209.
82. Jansson E, Terrados N, Norman B, Kaijser L: **Effects of training at simulated high altitude on exercise at sea level.** *Scand J Med Sci Sports* 1992, 2:2–6.
83. Terrados N: **Altitude training and muscular metabolism.** *Int J Sports Med* 1992, 13:S206–S209.
84. Green HJ: **Adaptations at extreme altitude: metabolic implications during exercise.** *Int J Sports Med* 1992, 13(Suppl 1):S163–S165.
85. Terrados N, Jansson E, Sylven C, Kaijser L: **Is hypoxia a stimulus for synthesis of oxidative enzymes and myoglobin?** *J Appl Physiol* 1990, 68:2369–2372.
86. Hoppeler H, Vogt M: **Muscle tissue adaptations to hypoxia.** *J Exp Biol* 2001, 204:3133–3139.
87. Rejnafarje C, Lozano R, Valdivieso J: **The polycythemia of high altitudes: iron metabolism and related aspects.** *Blood* 1959, 14:433–455.
88. Green HJ, Sutton JR, Cymerman A, et al.: **Operation Everest II: adaptations in human skeletal muscle.** *J Appl Physiol* 1989, 66:2454–2461.
89. Hoppeler H, Kleinert E, Schegel C, et al.: **Muscular exercise at high altitude: II morphological adaptation of skeletal muscle to chronic hypoxia.** *Int J Sport Med* 1990, 11:S3–S9.